

AD-A177 693

THE ACTIATION OF C-H BONDS WITH INTERMEDIATES PRODUCED  
BY MILD THERMAL A. (U) CALIFORNIA UNIV LOS ANGELES DEPT  
OF CHEMISTRY AND BIOCHEMISTR. P A CHETCUTI ET AL

1/1


UNCLASSIFIED

DEC 86 TR-126 N00014-86-K-0282

F/G 7/3

NL





MICROCOPY RESOLUTION TEST CHART

12

OFFICE OF NAVAL RESEARCH

Contract N00014-86-K-0282

TECHNICAL REPORT NO. 126

The Activation of C-H Bonds with Intermediates  
Produced by Mild Thermal and Photochemical Decomposition  
of the Metallacycle  $(C_5Me_5)Ir[C(p-ClC_6H_4)=NOC(=O)](CO)$   
in Hydrocarbon Solvents

by

Peter A. Chetcuti and M. Frederick Hawthorne\*

Prepared for Publication

in

*Journal of the American Chemical Society*

University of California at Los Angeles  
Department of Chemistry and Biochemistry  
Los Angeles, California 90024

December, 1986

Reproduction in whole or in part is permitted for  
any purpose of the United States Government

This document has been approved for public release and sale;  
its distribution is unlimited

DTIC  
ELECTE  
MAR 02 1987  
S D E

AD-A177 693

DTIC FILE COPY

87 3 2 013

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE

## REPORT DOCUMENTATION PAGE

1a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED			1b. RESTRICTIVE MARKINGS	
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION / AVAILABILITY OF REPORT  Distribution list enclosed	
2b. DECLASSIFICATION / DOWNGRADING SCHEDULE				
4. PERFORMING ORGANIZATION REPORT NUMBER(S)  126			5. MONITORING ORGANIZATION REPORT NUMBER(S)	
6a. NAME OF PERFORMING ORGANIZATION UCLA Dept. of Chem. & Biochem.		6b. OFFICE SYMBOL (if applicable)	7a. NAME OF MONITORING ORGANIZATION  Office of Naval Research/Chemistry Division	
6c. ADDRESS (City, State, and ZIP Code) 6115 Young Hall 405 Hilgard Avenue Los Angeles, CA 90024			7b. ADDRESS (City, State, and ZIP Code)  Department of the Navy Arlington, VA 22217	
8a. NAME OF FUNDING / SPONSORING ORGANIZATION Office of Naval Research		8b. OFFICE SYMBOL (if applicable)	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER  N00014-86-K-0282/RNT Code: 4135004	
8c. ADDRESS (City, State, and ZIP Code)			10. SOURCE OF FUNDING NUMBERS	
			PROGRAM ELEMENT NO 61153N	PROJECT NO.
			TASK NO.	WORK UNIT ACCESSION NO.
11. TITLE (Include Security Classification) The Activation of C-H Bonds with Intermediates Produced by Mild Thermal and Photochemical Decomposition of the Metallacycle $(C_5Me_5)Ir[C(p-ClC_6H_4)=NOC(=O)](CO)$ in Hydrocarbon Solvents				
12. PERSONAL AUTHOR(S) Peter A. Chetcuti and M. Frederick Hawthorne*				
13a. TYPE OF REPORT Technical Report		13b. TIME COVERED FROM TO		14. DATE OF REPORT (Year, Month, Day) 02/18/87
15. PAGE COUNT 1				
16. SUPPLEMENTARY NOTATION To be published in Journal of the American Chemical Society				
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)	
FIELD	GROUP	SUB-GROUP	Metallacycle; Iridacycle; C-H Bond Activation; Nitriles Side-bonded Nitrile Complex; ←	
19. ABSTRACT (Continue on reverse if necessary and identify by block number) Thermolysis of the metallacycle $(\eta^5-C_5Me_5)Ir[C(p-ClC_6H_4)=NOC(=O)](CO)$ (1) at 50°C in benzene or cyclohexane leads initially to the formation of the side-bonded nitrile complex $(\eta^5-C_5Me_5)Ir(\eta^2-NCC_6H_4)(CO)$ (5) which has been isolated and characterized. Upon further heating at 50°C loss of the nitrile ligand from 5 results in C-H bond activation of benzene solvent to generate $(\eta^5-C_5Me_5)Ir(H)(C_6H_5)(CO)$ (6) in 90% yield. These reactions presumably proceed through the 16-electron metal fragment " $(\eta^5-C_5Me_5)Ir(CO)$ ". Decomposition of 1 and 5 under mild photolytic conditions in both benzene and cyclohexane also led to C-H activation of the hydrocarbon solvents, generating 2 and $(\eta^5-C_5Me_5)Ir(CO)(H)(C_6H_{11})$ , 6a.				
Keywords: <i>Key words:</i>				
20. DISTRIBUTION / AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT <input type="checkbox"/> DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION Unclassified	
22a. NAME OF RESPONSIBLE INDIVIDUAL M. F. Hawthorne			22b. TELEPHONE (Include Area Code) (213) 825-7376	22c. OFFICE SYMBOL

The Activation of C-H Bonds with Intermediates Produced by  
Mild Thermal and Photochemical Decomposition of the  
Metallacycle  $(\eta^5\text{-C}_5\text{Me}_5)\text{Ir}[\text{C}(\text{p-ClC}_6\text{H}_4)=\text{NOC}(=\text{O}))(\text{CO})$   
in Hydrocarbon Solvents

Peter A. Chetcuti and M. Frederick Hawthorne\*

Department of Chemistry and Biochemistry  
University of California, Los Angeles  
Los Angeles, California 90024



Accession For	
NTIS	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution	
Availability Codes	
Dist	
A-1	

Abstract

Thermolysis of the metallacycle  $(\eta^5\text{-C}_5\text{Me}_5)\text{Ir}[\text{C}(\text{p-ClC}_6\text{H}_4)=\text{NOC}(=\text{O}))(\text{CO})$  (**1**) at 50°C in benzene or cyclohexane leads initially to the formation of the side-bonded nitrile complex  $(\eta^5\text{-C}_5\text{Me}_5)\text{Ir}(\eta^2\text{-NCC}_6\text{H}_4\text{Cl})(\text{CO})$  (**5**) which has been isolated and characterized. Upon further heating at 50°C loss of the nitrile ligand from **5** results in C-H bond activation of benzene solvent to generate  $(\eta^5\text{-C}_5\text{Me}_5)\text{Ir}(\text{H})(\text{C}_6\text{H}_5)(\text{CO})$  (**2**) in 90% yield. These reactions presumably proceed through the 16-electron metal fragment " $(\eta^5\text{-C}_5\text{Me}_5)\text{Ir}(\text{CO})$ ". Decomposition of **1** and **5** under mild photolytic conditions in both benzene and cyclohexane also led to C-H activation of the hydrocarbon solvents, generating **2** and  $(\eta^5\text{-C}_5\text{Me}_5)\text{Ir}(\text{CO})(\text{H})(\text{C}_6\text{H}_{11})$ , **6**.

The Activation of C-H Bonds with Intermediates Produced by  
Mild Thermal and Photochemical Decomposition of the  
Metallacycle  $(\eta^5\text{-C}_5\text{Me}_5)\text{Ir}[\text{C}(\text{p-ClC}_6\text{H}_4)=\text{NOC}(=\text{O})](\text{CO})$   
in Hydrocarbon Solvents

Peter A. Chetcuti and M. Frederick Hawthorne\*

Department of Chemistry and Biochemistry  
University of California, Los Angeles  
Los Angeles, California 90024

In recent years a number of transition metal complexes have been found to undergo photochemically induced intermolecular C-H bond activation of their hydrocarbon solvents.<sup>1-6</sup> The aryl and alkyl metal hydride products of these reactions are believed to result from the oxidative addition of the 16-electron coordinatively unsaturated metal fragments, "ML<sub>n</sub>", generated by the photolytically induced loss of H<sub>2</sub> or CO, into the C-H bonds of hydrocarbons (Scheme 1). Insertion into C-H bonds also occurs with thermal activation, suggesting that these reactions do not specifically require the reactive metal fragment to be in a photoexcited state.<sup>1a,4,7-18</sup> A preponderance of the previously reported examples of thermal C-H bond activation, in which the products of direct

[Scheme 1]

oxidative addition are observed, involve the reductive elimination of an alkyl or aryl metal hydride, to apparently produce the reactive species "ML<sub>n</sub>", followed by its oxidative addition into the C-H bond of another hydrocarbon.<sup>1b,4,7-9,15-18</sup> In effect, these reactions require as a reactant a metal center which has already undergone C-H activation into a C-H bond and constitute hydrocarbon exchange reactions.

We now wish to report the thermal activation of C-H bonds of hydrocarbons using the precursor metallacycle  $(\eta^5\text{-C}_5\text{Me}_5)\text{Ir}[\text{C}(p\text{-ClC}_6\text{H}_4)=\text{NOC}(=\text{O}))(\text{CO})$  (**1**).<sup>19,20</sup> Thermolysis of **1** would be expected to generate the 16-electron four-coordinate species " $(\eta^5\text{-C}_5\text{Me}_5)\text{Ir}(\text{CO})$ " (Scheme 2)

[Scheme 2]

which is presumed to be responsible for the activation of a number of hydrocarbons when generated photolytically from  $(\eta^5\text{-C}_5\text{Me}_5)\text{Ir}(\text{CO})_2$ .<sup>2</sup> When the metallacycle **1** was maintained in the dark at 50°C in benzene solution for four weeks, a hydride resonance at -15.07 ppm was observed and attributed to the product  $(\eta^5\text{-C}_5\text{Me}_5)\text{Ir}(\text{CO})(\text{H})(\text{C}_6\text{H}_5)$  **2a** (**2**) formed by the oxidative addition of " $(\eta^5\text{-C}_5\text{Me}_5)\text{Ir}(\text{CO})$ " into the C-H bonds of benzene;<sup>21</sup> addition of  $\text{CCl}_4$  converted **2** to the known  $(\eta^5\text{-C}_5\text{Me}_5)\text{Ir}(\text{CO})(\text{Cl})(\text{C}_6\text{H}_5)$  **22,2a** (**3**) in 90.4% overall yield while the known dimer  $[(\eta^5\text{-C}_5\text{Me}_5)\text{Ir}(\mu\text{-CO})]_2$  (**4**),<sup>23</sup> identified by its  $^1\text{H}$  NMR resonance at 1.58 ppm in  $\text{C}_6\text{D}_6$ , was produced in 8.6% yield as determined by NMR. When the reaction of **1** was carried out in 100% deuterated benzene and monitored by  $^1\text{H}$  NMR, two sets of doublets at  $\delta$  7.71 ( $J = 8.3$  Hz) and 7.01 ( $J = 8.3$  Hz) appeared;<sup>24</sup> as the reaction proceeded these resonances decreased in intensity as the resonances due to the free  $p\text{-ClC}_6\text{H}_4\text{CN}$  increased. The same intermediate resonances were observed when the reaction was carried out in cyclohexane and the  $^1\text{H}$  NMR of the reaction products obtained before the reaction had gone to completion, thus suggesting the existence of a common intermediate which is formed before oxidative addition of the solvent occurs. In order to isolate the intermediate, a THF solution of (**1**) was stirred at 50°C for 14 hours in the dark. Upon removal of the THF under *vacuo* and extraction of the residue with pentane, a yellow solution was obtained which on cooling afforded a yellow crystalline product in 66% yield. The complex obtained is a side-bonded nitrile complex  $(\eta^5\text{-C}_5\text{Me}_5)\text{Ir}(\eta^2\text{-NCC}_6\text{H}_4\text{Cl})\text{CO}$  **24** (**5**) identified by its characteristic low CN stretching frequency observed at  $1781\text{ cm}^{-1}$  in the IR spectrum.<sup>25,26</sup> Complex **5** was also characterized by  $^1\text{H}$  NMR spectroscopy, complete elemental analyses and its molecular structure has been determined by a single-crystal X-ray diffraction study.<sup>27</sup> The resonances in the  $^1\text{H}$  NMR spectrum of **5** **24** coincide with those of the intermediate observed when the thermolysis of **1** was monitored by  $^1\text{H}$  NMR as described above. In order to confirm that **5** independently activates C-H bonds of hydrocarbons, a 0.07 M solution of **5** in benzene was

maintained in the dark at 50°C for 36 hours, generating the phenyl hydride product, **2**, and the dimer, **4**, in 76.2% and 18.4% yield, respectively. A mechanism consistent with these observations involves the loss of CO<sub>2</sub> from **1** to generate the nitrile complex **5** which at 50°C is labile and loses the coordinated nitrile to release the metal fragment "(C<sub>5</sub>Me<sub>5</sub>)Ir(CO)" which then undergoes C-H bond activation (Scheme 3). The metallacycle **1** was also found to be very photosensitive; when a

[Scheme 3]

sample of **1** was maintained in benzene at 50°C and photolyzed<sup>28</sup> for 36 hours, **2** was obtained in 75.4% yield and **4** in 7.5% yield. The side-bonded nitrile **5** was observed when the <sup>1</sup>H NMR spectrum of the reaction mixture was obtained before the reaction had gone to completion although the steady state concentration of **5** was low since this compound was particularly photosensitive. When **5** was photolyzed<sup>28</sup> in benzene for 12 hours at 50°C, **2** was obtained in 80.6% yield by <sup>1</sup>H NMR; compound **4** was obtained in 4.7% yield. Photolysis of metallacycle **1** in cyclohexane for 13 days under the same conditions yielded the cyclohexylhydride product, (η<sup>5</sup>-C<sub>5</sub>Me<sub>5</sub>)Ir(CO)-(H)(C<sub>6</sub>H<sub>11</sub>) (**6**); addition of CCl<sub>4</sub> converted **6** to the known (η<sup>5</sup>-C<sub>5</sub>Me<sub>5</sub>)Ir(CO)(Cl)(C<sub>6</sub>H<sub>11</sub>) (**7**) **2a** in 48.6% overall yield. The dimer **4** was obtained in 33.7% yield. When the side-bonded nitrile complex **5** was photolyzed for 24 hours in cyclohexane at 50°C, **6** was obtained which was converted to the chloride derivative to yield **7** in 67.1% overall yield; compound **4** was obtained in 13.1% yield.

The rhodium metallacycle (η<sup>5</sup>-C<sub>5</sub>Me<sub>5</sub>)(PMe<sub>3</sub>)Rh[ $\overbrace{C(p\text{-FC}_6\text{H}_4)=\text{NOC}(=\text{O})}$ ] demonstrates a reactivity similar to that of **1** since the formation of a side-bonded nitrile complex and thermal C-H activation of hydrocarbon solvents are observed. Full details concerning this reaction will be reported shortly.


Acknowledgment. The authors are grateful to the Office of Naval Research for support of this research and the Johnson-Matthey Corporation for a generous gift of iridium and rhodium chloride. We would also like to thank Professor W. A. G. Graham for helpful discussions.

## References


1. a) Janowicz, A. H.; Bergman, R. G. *J. Am. Chem. Soc.* **1982**, *104*, 352. b) Janowicz, A. H.; Bergman, R. G. *J. Am. Chem. Soc.* **1983**, *105*, 3929. c) Wenzel, T. J.; Bergman, R. G. *J. Am. Chem. Soc.* **1986**, *108*, 4856.
2. a) Hoyano, A. K.; Graham, W. A. G. *J. Am. Chem. Soc.* **1982**, *104*, 3723. b) Hoyano, A. K.; McMaster, A. D.; Graham, W. A. G. *J. Am. Chem. Soc.* **1983**, *105*, 7190.
3. Prelana, R. A.; Bergman, R. G. *Organometallics* **1984**, *3*, 508.
4. Jones, W. D.; Feher, F. J. *J. Am. Chem. Soc.* **1984**, *106*, 1650.
5. Berry, M.; Elmitt, K.; Green, M. L. H. *J. Chem. Soc., Dalton Trans.* **1979**, 1950.
6. Rausch, M. D.; Gastinger, R. G.; Gardner, S. A.; Brown, R. K.; Wood, J. S. *J. Am. Chem. Soc.* **1977**, *99*, 7870.
7. Wax, M. J.; Stryker, J. M.; Buchanan, J. M.; Kovac, C. A.; Bergman, R. G. *J. Am. Chem. Soc.* **1984**, *106*, 1121.
8. Cooper, N. J.; Green, M. L. H.; Mahtab, R. *J. Chem. Soc., Dalton Trans.* **1979**, 1557.
9. Watson, P. A. *J. Chem. Soc., Chem. Commun.* **1983**, 276.
10. Watson, P. A. *J. Am. Chem. Soc.* **1983**, *105*, 6491.
11. Baudry, D.; Ephritikhine, M.; Felkin, H. *J. Chem. Soc., Chem. Commun.* **1980**, 1243.
12. Crabtree, R. H.; Mihelcic, J. M.; Quirk, J. M. *J. Am. Chem. Soc.* **1979**, *101*, 7738.
13. Crabtree, R. H.; Mellea, M. F.; Mihelcic, J. M.; Quirk, J. M. *J. Am. Chem. Soc.* **1982**, *104*, 107.
14. Burk, M. J.; Crabtree, R. H.; Parnell, C. P.; Uriarte, R. J. *Organometallics* **1984**, *3*, 816.
15. Fendrick, C. M.; Marks, T. M. *J. Am. Chem. Soc.* **1986**, *108*, 425.
16. Bruno, J. W.; Smith, G. M.; Marks, T. J.; Fair, C. K.; Schultz, A. J.; Williams, J. M. *J. Am. Chem. Soc.* **1986**, *108*, 40.
17. Chatt, J.; Davidson, J. M. *J. Chem. Soc.* **1965**, 843.

18. a) Ittel, S. D.; Tolman, C. A.; English, A. D.; Jesson, J. P. *J. Am. Chem. Soc.* **1976**, *98*, 6073. b) Ittel, S. D.; Tolman, C. A.; English, A. D.; Jesson, J. P. *J. Am. Chem. Soc.* **1978**, *100*, 4080. c) Ittel, S. D.; Tolman, C. A.; English, A. D.; Jesson, J. P. *J. Am. Chem. Soc.* **1979**, *101*, 1742.
19. We have synthesized a number of metallacycles by cycloaddition of aryl nitrile oxides with low valent metal carbonyl complexes. A preliminary communication has been published (Walker, J. A.; Knobler, C. B.; Hawthorne, M. F. *J. Am. Chem. Soc.* **1983**, *105*, 3370) and a complete report will be submitted shortly.
20. Selected data for **1** (full details will be reported elsewhere<sup>19</sup>).  
**1**: Anal. Calcd. for  $C_{19}H_{19}ClIrNO$ : C, 42.57; H, 3.58; Ir, 35.85; N, 2.61. Found: C, 42.12; H, 3.64; Ir, 35.88; N, 2.40.  $^1H$  NMR ( $C_6D_6$ ):  $\delta$  7.39 (d, 2H,  $J = 8.5$  Hz), 7.10 (d, 2H,  $J = 8.4$  Hz), 1.23 (s, 15H).
21. The reactions were carried out in NMR tubes using dried and degassed solvents. In a typical experiment 0.8 mL of reactant solvent was vacuum transferred into 8 mg of the metallacycle **1** and the reactants maintained at a constant temperature. When the reaction was complete the reactant solvent was removed under vacuum and deuterated benzene was then vacuum transferred in and the tube sealed. The alkyl and aryl metal hydrides formed were identified by matching the  $^1H$  NMR chemical shifts with the reported values published by other workers or by conversion to the chloro derivatives by addition of  $CCl_4$ .<sup>2a</sup> Relative yields were measured by  $^1H$  NMR integration.
22. Kang, J. W.; Maitlis, P. M. *J. Organomet. Chem.* **1971**, *26*, 393.
23. Graham, W. A. G.; personal communication.
24. **5**: IR (nujol): 1949 (br,  $\nu_{CO}$ ), 1781 (m,  $\nu_{CN}$ )  $cm^{-1}$ .  $^1H$  NMR ( $C_6D_6$ ):  $\delta$  7.72 (d, 2H,  $J = 8.3$  Hz), 7.01 (d, 2H,  $J = 8.3$  Hz), 1.69 (s, 15H).
25. Storhoff, B. N.; Huntley, Jr., C. L. *Coord. Chem. Rev.* **1977**, *23*, 1.

26. Side-bonded nitrile complexes have been isolated and characterized as the major products in the thermolysis of the metallacycles  $(C_5H_5)Ir[C(Ar)=NOC(=O)](PPh_3)$ , ( $Ar = p-ClC_6H_4^-$ ,  $p-FC_6H_4^-$ ); Chetcuti, P. A.; Knobler, C. B.; Hawthorne, M. F. *Organometallics* **1986**, *5*, 1913.
27. Complete details on the X-ray crystal structure of **5** will be given in a full report discussing the reactivity of metallacycles of rhodium and iridium.
28. Photochemical experiments were carried out using a 100-W bulb which was supported six inches above an oil bath in which the NMR tubes were maintained at constant temperature. The radiated light was contained within the system by aluminum foil and a stream of cold nitrogen gas was used to cool the system to the desired temperature.



Scheme 1



Scheme 2

TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No. Copies</u>		<u>No. Copies</u>
Office of Naval Research Attn: Code 1113 800 N. Quincy Street Arlington, Virginia 22217-5000	2	Dr. David Young Code 334 NORDA NSTL, Mississippi 39529	1
Dr. Bernard Douda Naval Weapons Support Center Code 50C Crane, Indiana 47522-5050	1	Naval Weapons Center Attn: Dr. Ron Atkins Chemistry Division China Lake, California 93555	1
Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko, Code L52 Port Hueneme, California 93401	1	Scientific Advisor Commandant of the Marine Corps Code RD-1 Washington, D.C. 20380	1
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	12 high quality	U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 12211 Research Triangle Park, NC 27709	1
DTNSRDC Attn: Dr. H. Singerman Applied Chemistry Division Annapolis, Maryland 21401	1	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	1
Dr. William Tolles Superintendent Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375-5000	1	Naval Ocean Systems Center Attn: Dr. S. Yamamoto Marine Sciences Division San Diego, California 91232	1

END

4-87

DTIC